













SEXUAL EXPOSURE CHART

(if every person has only the same number of partners as you)

Number of Sexual Partners

Number of People Exposed To

1		1
2		3
3		7
4		15
5		31
6		63
7		127
8		255
9		511
10		1023
11		2047
12		4095

"When you have sex with someone, you are having sex with everyone they have had sex with for the last ten years, and everyone they and their partners have had sex with for the last ten years."

"Many teenagers, as well as adults, are indirectly exposed to more than one sexual partner each year because their partner has had sex with someone else."

The Most Common STD's

STD	SYMPTOMS	DAMAGE	NUMBERS
Chlamydia	Men- <u>Usually none.</u> Some experience discharge from the penis and burning with urination; Women- <u>Usually none.</u> Some experience pain and /or vaginal discharge.	Men- <u>Sterility.</u> Women- Considered to be the most common cause of sterility. Advanced stage may require removal of uterus, tubes and ovaries.	1) 4 million new cases reported annually. 2) 70% are not aware they are infected. 3) 40% of those sexually active are suspected carriers.
Gonorrhea	Men- Pus from urethra. Burning upon urination. Women- Pelvic pain. Painful urination and pus-like discharge.	Men- <u>Sterility, scarring of urethra and urinary tract problems.</u> Women- <u>Sterility, surgery in severe cases.</u>	1) 1.1 million new cases reported annually. 2) 80% are not aware they are infected in the early stages.
Syphilis	Men- 1st stage: swollen non-painful ulcers on external genitalia. 2nd stage: changes of skin, fever, enlarged lymph nodes. Women- 1st stage: <u>often unnoticed.</u> 2nd stage: same as men.	Men- <u>Brain disorders, heart disease, blindness, dementia and death.</u> Women- Same as men. Can cause birth defects or <u>death in newborns, if mother is infected.</u>	1) 120,000 new cases reported annually. 2) 50% are not aware they have this disease.
HIV/AIDS	Men- Fever, sore throat, fatigue, swollen lymph glands. Women- Same as men.	Men- Immune system breakdown, death. Women- Same as men.	1) 1 million infected. Approx. 20% in young adult population. 2) 20% infected during teens.
Genital Herpes	Men- Lesions appear at the site of infection- periodic eruptions of painful blister and ulcers anywhere on body. Women- Same as men.	Men- Continuous outbreaks. Eventually may lead to hospitalization. <u>Lifetime medication required.</u> Women- Same as men. Causes infant death.	1) 3.1 million infected. 500,000 new cases annually. 2) 75% show no symptoms. 3) 33% of women have this virus.
HPV Human Papilloma-Virus	Men- Wart-like genital growths. Women- (Often no visible symptoms.) Some experience vulva burning, itching and pain. Abnormal pap smears.	Men- Cancer of penis and anus. Women- Initially pre-malignant changes difficult to eradicate. Develops into cancer if not treated.	1) 24-40 million people are thought to be carriers. 2) 60 different strains exist 3) 33% of women have this virus
Hepatitis B	Men- <u>Often none.</u> Some experience yellowing of skin. (jaundice), tiredness, dark urine and gray-colored stool. Women- Same as men.	Men- <u>Severe liver damage.</u> Can eventually lead to cancer of the liver and cirrhosis. Women- Same as men.	1) 300,000 new cases annually 2) Most common STD in the world. 3) Approximately 1.25 million chronically infected Americans.
PID Pelvic-Inflammatory Disease	Men- This disease affects women only. Women- See Chlamydia & Gonorrhea. Often symptoms are minor and mistaken for menstrual cramps.	Men- The disease affects women only. Women- *Ectopic Pregnancy (fertilized egg implants in the fallopian tubes instead of uterus. (Can be <u>Life-Threatening.</u>) Sterility & cancer common.	1) 1 million new cases annually most in 15-19 year age group. 2) 100,000 women become infertile as a result of PID each year.